Clinician-Rated Dimensions of Psychosis Symptom Severity

Name:

Age:

Sex: [] Male [] Female

Date:

Instructions: Based on all the information you have on the individual and using your clinical judgment, please rate (with checkmark) the presence and severity of the following symptoms as experienced by the individual in the past seven (7) days.

Domain	0	1	2	3	4	Score
I. Hallucinations	□Not present	Equivocal (severity or duration not sufficient to be considered psychosis)	Present, but mild (little pressure to act upon voices, not very bothered by voices)	Present and moderate (some pressure to respond to voices, or is somewhat bothered by voices)	Present and severe (severe pressure to respond to voices, or is very bothered by voices)	
II. Delusions	□Not present	Equivocal (severity or duration not sufficient to be considered psychosis)	Present, but mild (little pressure to act upon delusional beliefs, not very bothered by beliefs)	Present and moderate (some pressure to act upon beliefs, or is somewhat bothered by beliefs)	□ Present and severe (severe pressure to act upon beliefs, or is very bothered by beliefs)	
III. Disorganized speech	□Not present	□ Equivocal (severity or duration not sufficient to be considered disorganization)	Present, but mild (some difficulty following speech)	Present and moderate (speech often difficult to follow)	Present and severe (speech almost impossible to follow)	
IV. Abnormal psychomotor behavior	□Not present	Equivocal (severity or duration not sufficient to be considered abnormal psychomotor behavior)	Present, but mild (occasional abnormal or bizarre motor behavior or catatonia)	Present and moderate (frequent abnormal or bizarre motor behavior or catatonia)	Present and severe (abnormal or bizarre motor behavior or catatonia almost constant)	
V. Negative symptoms (restricted emotional expression or avolition)	□Not present	Equivocal decrease in facial expressivity, prosody, gestures, or self-initiated behavior	Present, but mild decrease in facial expressivity, prosody, gestures, or self-initiated behavior	Present and moderate decrease in facial expressivity, prosody, gestures, or self-initiated behavior	Present and severe decrease in facial expressivity, prosody, gestures, or self-initiated behavior	
VI. Depression	□Not present	□ Equivocal (occasionally feels sad, down, depressed, or hopeless; concerned about having failed someone or at something but not preoccupied)	□ Present, but mild (frequent periods of feeling very sad, down, moderately depressed, or hopeless; concerned about having failed someone or at something, with some preoccupation)	□ Present and moderate (frequent periods of deep depression or hopelessness; preoccupation with guilt, having done wrong)	□ Present and severe (deeply depressed or hopeless daily; delusional guilt or unreasonable self-reproach grossly out of proportion to circumstances)	
VII. Mania	□Not present	Equivocal (occasional elevated, expansive, or irritable mood or some restlessness)	Present, but mild (frequent periods of somewhat elevated, expansive, or irritable mood or restlessness)	Present and moderate (frequent periods of extensively elevated, expansive, or irritable mood or restlessness)	□ Present and severe (daily and extensively elevated, expansive, or irritable mood or restlessness)	

Copyright © 2013 American Psychiatric Association. All Rights Reserved.

This material can be reproduced without permission by researchers and by clinicians for use with their patients